

MATH 15 MIDTERM SOLUTIONS
S'01, May 7, 2001

1. (5 pts) Find $f^{-1}(3)$, if

$$f(x) = \frac{x+5}{x+1}$$

$$\begin{aligned}\frac{x+5}{x+1} &= 3 \\ x+5 &= 3x+3 \\ 2 &= 2x \\ x &= 1\end{aligned}$$

So $f^{-1}(3) = 1$.

2. (5 pts) Find an equation of the line through $(3, 4)$ and $(-1, -2)$.

$$\text{slope} = \frac{4 - (-2)}{3 - (-1)} = \frac{6}{4} = \frac{3}{2}$$

So the equation is

$$\begin{aligned}\frac{y - (-2)}{x - (-1)} &= \frac{3}{2} \\ y + 2 &= \frac{3}{2}(x + 1) \\ y &= \frac{3}{2}x + \frac{3}{2} - 2 \\ y &= \frac{3}{2}x - \frac{1}{2}\end{aligned}$$

3. (5 pts each)
(a) Show that

$$2 - 3 \log 5 = \log \left(\frac{4}{5} \right)$$

Notice $2 = \log 100$, so

$$2 - 3 \log 5 = \log 100 - \log 5^3 = \log \left(\frac{100}{125} \right) = \log \left(\frac{4}{5} \right).$$

- (b) Let $a = \log x$ and $b = \log y$. Express in terms of a and b


$$\log \left(\frac{\sqrt[3]{x}}{\sqrt{y^7}} \right).$$

$$\log \left(\frac{\sqrt[3]{x}}{\sqrt{y^7}} \right) = \log \left(\frac{x^{1/3}}{(y^7)^{1/2}} \right) = \log x^{1/3} - \log y^{7/2} = \frac{\log x}{3} - \frac{7}{2} \log y = \frac{a}{3} - \frac{7b}{2}$$

4. (10 pts) Graph the function

$$f(x) = \begin{cases} x+2 & \text{if } x < -1 \\ -x & \text{if } -1 \leq x < 2 \\ x-2 & \text{if } 2 \leq x \end{cases}.$$

Find $f(f(3))$.



$$f(f(3)) = f(3 - 2) = f(1) = -1$$

5. (10 pts) Find the domain and the range of

$$f(x) = \frac{1}{\sqrt{x-1}}.$$

For the domain, you want $x-1 \geq 0$ because of the square root and $x-1 \neq 0$ because of the division. So $x-1 > 0$, that is $x > 1$ and $D(f) = \{x \mid x > 1\}$.

For the range, notice that $x-1$ can be any positive number, so $\sqrt{x-1}$ can be any positive number, so $\frac{1}{\sqrt{x-1}}$ can be any positive number too. Alternately:

$$\begin{aligned} \frac{1}{\sqrt{x-1}} &= y && \text{hence } y \geq 0 \\ \frac{1}{x-1} &= y^2 \\ 1 &= (x-1)y^2 && \text{shows } y \neq 0 \\ \frac{1}{y^2} &= x-1 \\ \frac{1}{y^2} + 1 &= x \end{aligned}$$

which gives some x for any $y \neq 0$. But we already know $y \geq 0$. Hence $R(f) = \{y \mid y > 0\}$.

6. (15 pts) A bacterium culture is growing exponentially. After 1 day, it weighs 60 g, after 3 days 135 g.

(a) Find a function $f(t)$ that describes the weight of the culture after t days.

We know $f(t) = ab^t$ for some a and b . Now

$$(1) \quad 60 = ab$$

$$(2) \quad 135 = ab^3$$

Divide (2) by (1) to get

$$\frac{135}{60} = \frac{ab^3}{ab} \implies \frac{9}{4} = b^2 \implies b = \frac{3}{2}$$

Now use (1) to find a :

$$\begin{aligned} 60 &= a \frac{3}{2} \\ 40 &= a \end{aligned}$$

So $f(t) = 40 \left(\frac{3}{2}\right)^t$.

(b) What was the weight of the bacteria initially?

$f(0) = 40$ so there were 40 g of bacteria initially.

(c) What is the percent increase per day?

$\frac{3}{2} = 150\%$, so the increase is 50%.

(d) Assuming the growth were linear at a rate of 50%/day, how much would the culture weigh after 3 days?

So it would grow by 50%(40 g) = 20 g on the first day, and the 20 g each day to 100 g after 3 days.